

## Cloning and Characterization of a New Isoform of Mouse Interleukin-18

Yong-Jie YANG, Zhao-Yuan WANG, Song-Hua CHEN, and Xi-Rui GE\*

*Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences;  
Graduate School of the Chinese Academy of Sciences, Shanghai 200031, China*

**Abstract** Interleukin-18 (IL-18) is a novel proinflammatory cytokine with potent interferon (IFN)- $\gamma$  inducing activity that plays an important biological role in the enhancement of the activity of natural killer cells and cytotoxic T lymphocytes. In this study, we have identified a novel short form of IL-18 in mouse, named IL-18s. IL-18s might be an alternative splicing variant of IL-18 and its cDNA contains a 57 bp in-frame deletion. Like IL-18, IL-18s is also widely expressed in mouse tissues. It was suggested that IL-18s might have a caspase-1-dependent mechanism for maturation and secretion similar to that of IL-18: when transfected in COS-7 cells, pro-IL-18s (22 kDa) could be detected, and the mature IL-18s (16 kDa) could also be detected when combined with caspase-1. We observed that recombinant mouse IL-18s did not show any IL-18-like function, and IL-18s could enhance the ability of IL-18 to increase IFN- $\gamma$  production by approximately 40% in mouse splenocytes. This effect was observed primarily at relative low concentrations of IL-18, suggesting that IL-18s might regulate the activity of IL-18 in the physiological conditions.

**Key words** IL-18; isoform; IL-18s; IFN- $\gamma$ ; caspase-1; factor Xa

Interleukin-18 (IL-18), originally identified as an interferon (IFN)- $\gamma$  inducing factor, was first isolated from the livers of mice stimulated with *Propionibacterium acnes* and lipopolysaccharide (LPS) [1]. IL-18 is a potent pro-inflammatory cytokine that induces IFN- $\gamma$  production in T cells and natural killer (NK) cells [2], enhances the Fas ligand and perforin-mediated T cell and NK cell cytotoxicity [3–8], and plays a critical role in the T-lymphocyte helper type 1 response [8]. IL-18 also induces expression of GM-CSF, inflammatory cytokines (e.g., tumour necrosis factor- $\alpha$ , IL-1 $\beta$  and IL-13), and chemokines, such as IL-8, macrophage inflammatory protein (MIP)-1 $\alpha$  and MIP-1 $\beta$  [2,10–12].

IL-18 is structurally related to IL-1 $\beta$  and has been shown to be a member of the IL-1 superfamily [13,14]. Furthermore, the IL-18R complex consists of a ligand-binding subunit (IL-18R $\alpha$ ) and a signaling subunit (IL-18R $\beta$ ), both of which are members of the IL-1R family [15–17].

Like IL-1 $\beta$ , IL-18 is synthesized in its precursor form, proIL-18, which requires caspase-1 cleavage to become a biologically active molecule [18–20]. However, unlike IL-1 $\beta$ , *IL-18* mRNA is expressed constitutively in many types of cells and IL-18 protein is stored in the producing cells as a biologically inactive precursor, which is partly due to the particular structure and characteristics of the *IL-18* gene [21–24]. LPS can induce the upregulation of *IL-18* mRNA and the secretion of mature IL-18 by the activation of caspase-1 [20–22]. In addition, the activation and secretion of IL-18 are regulated by caspase-1-independent mechanisms [18,21,25].

Recently, proIL-18 was observed combining with IgM in human blood plasma [26], however its physiological function is still not clear. Furthermore, a 51 bp in-frame deletion in the leading sequence of *IL-18* was also reported in rainbow trout [31]. *IL-18s* cDNA has been reported in rats with a 57 bp in-frame deletion [32], but further reports are not available.

In our study, a mouse *IL-18* isoform named *IL-18s*, 57 bp shorter than the normal *IL-18*, was cloned. We found that the *IL-18s* mRNA was constitutively expressed in a

Received: July 14, 2005 Accepted: September 29, 2005

This work was supported by a grant from the Pilot Project of Knowledge Innovation Program of the Chinese Academy of Sciences (No. KSCX 2-3-04-03)

\*Corresponding author: Tel, 86-21-54920407; Fax, 86-21-54920406; Email, gexirui@sibs.ac.cn

DOI: 10.1111/j.1745-7270.2005.00116.x

variety of tissues accompanying *IL-18*. The biochemical properties of IL-18s were further studied to find that IL-18s had a similar mechanism for maturation and secretion from the mammalian cells, which was caspase-1-dependent. The cloned *IL-18s* cDNA was then expressed in *Escherichia coli*, and recombinant mouse IL-18s did not exhibit any IL-18-like function. When IL-18 and IL-18s were both added to mouse splenocytes, we observed that IL-18s could modestly enhance the ability of IL-18 to induce IFN- $\gamma$ .

## Materials and Methods

### Mice and cell lines

BALB/c female mice 6–8 weeks of age were obtained from the Animal Center of the Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences (Shanghai, China). The COS-7 cell line was provided by the Cell Bank of Type Culture Collection, Chinese Academy of Sciences. COS-7 cells were cultured in Dulbecco's modified Eagle's medium and fresh isolated splenocytes were cultured in RPMI 1640 medium in a humidified atmosphere with 5% CO<sub>2</sub> at 37 °C. Both media were supplemented with 10% fetal bovine serum, 2 mM *L*-glutamine, 50 U/ml penicillin and 50  $\mu$ g/ml streptomycin.

### Reagents

LPS (a phenol-extracted preparation from *E. coli* 055: B5) was obtained from Sigma (St. Louis, USA), dissolved in water at 1 mg/ml, and stored at –20 °C. DOTAP

liposomal transfection reagent and protease inhibitor cocktails were purchased from Roche (Mannheim, Germany). Caspase inhibitor I (Z-VAD-FMK) was purchased from Calbiochem (San Diego, USA). Mouse IL-18 (pep-IL-18) and polyclonal rabbit antimurine IL-18 antibody were provided by PeproTech (Rocky Hill, USA). The enzyme-linked immunosorbent assay (ELISA) kit for mouse IFN- $\gamma$  was purchased from R&D Systems (Minneapolis, USA). Gel was analyzed by a gel imaging system (FR-200; Furi Tech, Shanghai, China).

### Reverse transcription-polymerase chain reaction (RT-PCR)

Specific primers (**Table 1**) were synthesized to amplify the full-length cDNA sequences of IL-18 and caspase-1. Total RNA was extracted from spleen of healthy BALB/c mouse using Trizol (Invitrogen, San Diego, USA) according to the manufacturer's instructions and reverse-transcribed into cDNA using Moloney murine leukemia virus reverse transcriptase (Promega, Madison, USA) in the presence of oligo(dT)<sub>15–18</sub>. All products amplified by PCR were directly cloned into pCR4TOPO (Invitrogen) for sequencing.

### IL-18s expression in mouse

To determine tissue distribution of *IL-18s* mRNA in healthy mouse, total RNA was extracted from organs including brain, small intestine, heart, liver, spleen, kidney, lung and pancreas. RT-PCR was performed using the primers 18q1/18q2, specifically detecting IL-18 and IL-18s expression. The PCR reaction programs were as follows: 94 °C for 4 min; 94 °C for 25 s, 58 °C for 20 s, 72 °C for 20 s,

**Table 1** Primers used to amplify full-length cDNA sequences of interleukin (IL)-18 and caspase-1

Point	Primer	Sequence (5'→3')
IL-18 cloning [9]	18C1 (upstream)	ACCTTCCAATCACTTCCTC
	18C2 (downstream)	CAGGCGAGGTCATCACAAGG
Caspase-1 cloning	ICE-1 (upstream)	CTGCGGTGTAGAAAAGAAACG
	ICE-2 (downstream)	GGCACGATTCTCAGCATAGG
IL-18 detection	18q1 (upstream)	GGCTGTGACCCTCTCTGTGAA
	18q2 (downstream)	TGGCAAGCAAGAAAGTGTCCCT
IL-18s propiece cloning	18sT1 (upstream)	AAC <b>GGATCC</b> ATGGCTGCCATGTCA
	18T2 (downstream)	AAAGTT <b>ACGTCCTTCGAT</b> GTCTCCATT
IL-18 propiece cloning	18T1 (upstream)	AAC <b>GAATTC</b> ATGGCTGCCATGTCA
	18T2 (downstream)	AAAGTT <b>ACGTCCTTCGAT</b> GTCTCCATT
Mature piece cloning	18T3 (upstream)	GGAGACATCGA <b>AGGACGTA</b> ACTTTGGC
	18T4 (downstream)	AAC <b>CTCGAGCTAACTT</b> GATGTAA

Restriction sites are shown in bold and factor Xa site mutation sequences are underlined.

30–35 cycles; and a final extension at 72 °C for 10 min. To detect IL-18s protein in mouse, mouse splenocytes were obtained as described previously [27] and lysed in buffer containing protease inhibitors and caspase inhibitor. The lysate was immunoprecipitated with rabbit antimurine IL-18 antibody and resolved by 12%–15% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) under reducing conditions. The gel was transferred to nitrocellulose membrane and incubated first with primary antibodies (1:1000) at 4 °C for 12–16 h, then with peroxidase-conjugated mouse anti-rabbit IgG. The membrane was developed by enhanced chemiluminescence.

### Construction of eukaryotic expression plasmids and transient expression of IL-18s

The full-length murine *proIL-18s*, *proIL-18* or *caspase-1* cDNA was subcloned into pcDNA3 with C-end myc-tag (pcDNA3-*proIL-18s*, pcDNA3-*proIL-18*, pcDNA3-*caspase-1*) by introducing the *Bam*HI (or *Eco*RI for *proIL-18*)/*Xho*I site at the 5' or 3' end. All the constructs were confirmed by DNA sequencing.

COS-7 cells were transfected with pcDNA3-*proIL-18s* (or pcDNA3-*proIL-18*) alone or in combination with pcDNA3-*caspase-1* in 35 mm dishes by DOTAP transfection according to the manual. In brief, 2 µg of DNA was diluted to a concentration of 0.1 µg/µl in HBS buffer (20 mM HEPES, 150 mM NaCl, pH 7.4); and 15 µl DOTAP reagent was diluted to a final volume of 50 µl with HBS. Then the DNA solution was gently mixed with the DOTAP solution and incubated for 10–15 min at room temperature. The COS-7 cells were washed with serum-free Opti-MEM1 (Gibco BRL, Gaithersburg, USA), plated in 1 ml Opti-MEM1 containing the DOTAP/DNA mixture for 6 h, then washed and incubated with fresh Opti-MEM1 medium supplemented with 50 U/ml penicillin and 50 µg/ml streptomycin. After 24 h the cells were fixed with 4% paraformaldehyde, and stained with mouse anti-myc monoclonal antibody and fluorescein-isothiocyanate (FITC)-labeled secondary antibody for immunofluorescence detection. Finally, the nuclei were stained with 4',6'-diamidino-2-phenylindole dihydrochloride (DAPI). The fluorescence of FITC and DAPI were observed using a confocal microscope. For Western blot detection, COS-7 cells were directly lysed in loading buffer or immunoprecipitation buffer containing protease inhibitors followed by immunoprecipitation with anti-IL-18 antibody 24 or 48 h post-transfection. The culture supernatants were collected and concentrated 200-fold by ultrafiltration (Centricon-10; Amicon, Beverly, USA) or concentrated 10-fold by immunoprecipitation with anti-IL-18 antibody. Then the supernatants, immunoprecipitates

and cell lysates were detected by Western blot.

### Construction of prokaryotic expression plasmids

The murine *proIL-18s* and *proIL-18* cDNA was used for generating a mutation of the caspase-1 cleavage site into a factor Xa site. Introduction of mutation in *proIL-18s* was performed by a two-step PCR reaction [28,29]. The propiece of IL-18s cDNA (or IL-18) was generated using primers 18sT1/18T2 (or 18T1/18T2 for IL-18) containing the *Bam*HI site (or *Eco*RI for *IL-18*) before the open reading frame (ORF). The mature piece of *IL-18s* or *IL-18* cDNA was generated by primers 18T3/18T4 containing the *Xho*I site after the ORF. The 123 bp propiece and 446 bp mature piece (503 bp for *IL-18*) of the *IL-18s* cDNA (or *IL-18* cDNA) were resolved by electrophoresis in 1% agarose and purified. These two cDNA pieces were mixed at 1:1 and used as the template for the second PCR step to generate a complete murine *IL-18s* cDNA in which the caspase-1 site is mutated into the factor Xa site. The IL-18s, IL-18s (Xa), and IL-18(Xa) cDNAs were ligated into the pGEX-4T-1 to form pGEX-*proIL-18s*, pGEX-*proIL-18s*(Xa) and pGEX-*proIL-18*(Xa), respectively, with the use of *Bam*HI/*Xho*I (for IL-18s) or *Eco*RI/*Xho*I (for IL-18). All the constructs were confirmed by DNA sequencing.

### Protein expression and purification

pGEX-*proIL-18s*, pGEX-*proIL-18s*(Xa) or pGEX-*proIL-18*(Xa) was transformed into *E. coli* BL21(DE3) competent cells and expressed as described previously [29,30]. In brief, 10 ml overnight culture from a fresh single colony of BL21 was added into 490 ml of 2×YT medium containing 100 µg/ml ampicillin and grown at 37 °C until its  $A_{600}$  reached 0.6. Protein expression was induced by isopropyl β-D-thiogalactopyranoside (IPTG; a final concentration of 0.1 mM). Then the culture was further incubated with shaking at 25 °C for 6 h. For preparation of the fusion proteins, bacteria were harvested by centrifugation (3500 g for 10 min at 4 °C) and rapidly resuspended in buffer A (20 mM Tris-HCl, pH 7.5, 5 mM EDTA, 0.1% Triton X-100). Lysozyme was added to a final concentration of 1 mg/ml. Cells were lysed on ice by mild sonication and the mixture was kept on ice for 15–20 min. The soluble proteins were obtained by centrifugation at 20,000 g, 4 °C for 20 min, mixed with 1 ml glutathione-Sepharose 4B beads (Amersham Pharmacia Biotech, Uppsala, Sweden), and incubated with mild shaking at 4 °C for 1 h. Then the beads were pelleted by centrifugation (500 g for 5 min), washed twice with 10 volumes of buffer A, twice with 10 volumes of buffer B (buffer A containing 300 mM NaCl), and twice with 10 volumes of cleavage

buffer (50 mM Tris-HCl, pH 8.0, 200 mM NaCl, 5 mM CaCl<sub>2</sub>).

### Cleavage of the fusion protein and purification of murine IL-18s

The fusion protein GST-proIL-18s, GST-proIL-18s(Xa) or GST-proIL-18(Xa) was incubated (22 °C, 3–12 h) with the appropriate amount of thrombin protease (for proIL-18s) or factor Xa (for mature IL-18s and IL-18) in 1 ml cleavage buffer according to the manual (Amersham Pharmacia Biotech) with constant shaking on a rotating wheel. Once the digestion was completed, the supernatant was collected by centrifugation (500 g for 5 min at 4 °C), and washed three times with 1 ml ice-cold phosphate-buffered saline. All the eluates were mixed, and clarified by centrifugation (12,000 g, 10 min) and ultrafiltration (Centricon-10). The purified protein concentration was measured using Bradford reagent (Bio-Rad, Hercules, USA) and stored at –70 °C. The fusion proteins before digestion and the concentrated elutes were resolved by 12%–15% SDS-PAGE followed by Coomassie blue staining.


### Bioassay of murine IL-18s

Mouse spleen cells were used to assay murine IL-18 as described previously [27]. Spleen cells were washed twice and cultured in RPMI 1640 medium in 48-well culture plates. Murine proIL-18s or mature IL-18s was added alone or in combination with pep-IL-18. Spleen cells were stimulated with 0.5 µg/ml LPS. The culture was incubated for 24 h at 37 °C and the supernatants were then assayed for mIFN-γ by ELISA.

## Results

### Molecular cloning of a novel mouse *IL-18* isoform

When cloning the mouse *IL-18* cDNA by RT-PCR from spleen cells, two distinct products were detected [Fig. 1(A)]. Subsequent DNA sequencing identified these two cDNAs: one is identical to the *IL-18* cDNA, and the other contains a 57 bp in-frame deletion that results in a 19 amino acid deletion in the C-terminal portion of the putative protein, accordingly named IL-18s. The 57 bp deletion, corresponding to nucleotide 355–411 of the full-length *IL-18* cDNA, occurs at the 5' end of exon 7 of the *IL-18* gene [Fig. 1(B)]. For the reconfirmation of the existence of *IL-18s*, a specific primer pair 18q1/18q2 was used and a similar result (a strong band for *IL-18*, and a faint band for *IL-18s*) was reproduced by RT-PCR [Fig. 1(C)].



**Fig. 1** Sequence of a novel short form of interleukin-18 (*IL-18s*) cDNA and confirmation of the deletion in *IL-18s* by reverse transcription-polymerase chain reaction (RT-PCR)


(A) Cloning of *IL-18* and *IL-18s*. The RNA was extracted from mouse splenocytes, and *IL-18* and *IL-18s* were amplified by RT-PCR. 1, 100 bp DNA ladder marker; 2, the full length of *IL-18* (676 bp) and *IL-18s* cDNA (619 bp); 3 and 4, the full-length cDNA of *IL-18* and *IL-18s*, respectively. (B) *IL-18* and *IL-18s* cDNA nucleotide sequences. Nucleotide positions are numbered on the left of the cDNA sequence. The deleted sequence is shown in bold. (C) Confirmation of the presence of the deletion in *IL-18s* by RT-PCR. The RNA was extracted from mouse splenocytes and specific primer pair (18q1/18q2) was used for the amplification of *IL-18* and *IL-18s*. 1, molecular mass marker; 2, negative control; 3, the PCR products of *IL-18* (201 bp) and *IL-18s* (144 bp).

### Expression of *IL-18s* mRNA and protein in mouse

As shown in Fig. 2(A), both *IL-18* and *IL-18s* were expressed in all the selected organs with a similar expression pattern detected previously in spleen [Fig. 1(C)], and a comparatively high expression of IL-18s was observed in pancreas. Similarly, two protein bands were observed in the lysates of fresh isolated splenocytes. The major band (approximately 24 kDa) corresponds to proIL-18, and the minor band (approximately 21–22 kDa), consistent with the predictive size, might correspond to proIL-18s [Fig. 2(B)].

### Expression of IL-18s in COS-7 cells

To determine the expression and distribution of IL-18 and IL-18s, we investigated the subcellular localization of




**Fig. 2** Expression of a novel short form of interleukin-18 (*IL-18s*) in mouse

(A) Tissue distribution of *IL-18s* mRNA expression. 1, heart; 2, liver; 3, lung; 4, kidney; 5, brain; 6, small intestine; 7, spleen; 8, pancreas; 9, negative control; 10, 100 bp DNA ladder marker. (B) The putative expression of proIL-18s protein in splenocyte. The cell lysates from freshly isolated splenocytes were immunoprecipitated with or without anti-IL-18 antibody (lanes 1 and 2, respectively). The immunoprecipitates were resolved by 12%–15% sodium dodecyl sulphate-polyacrylamide gel electrophoresis, and electrotransferred onto nitrocellulose membranes. The membranes were then probed with antibodies against IL-18.

them in COS-7 cells transfected with pcDNA3-IL-18 and pcDNA3-IL-18s observed by confocal microscopy. IL-18 was clearly detected in the cell cytoplasm only in the transfected cells and IL-18s localized similarly in the cytoplasm [Fig. 3(A)]. Both IL-18 and IL-18s showed as uneven dotted-distribution in the cytoplasm.

IL-18s shares the same leader sequence and potential caspase-1 cleavage site with IL-18. Thus we wanted to determine whether caspase-1 cleaves proIL-18s and facilitates the secretion of mature IL-18s. COS-7 cells were transfected with pcDNA3-IL-18s alone or in combination with pcDNA3-caspase-1. pcDNA3-IL-18 was used as a positive control in parallel. Twenty-four hours after transfection, the cells were directly lysed. The subsequent immunoblotting results revealed that a polypeptide of approximately 22 kDa corresponding to proIL-18s was recognized by anti-murine IL-18 antibody, and a 24 kDa band corresponding to proIL-18 was also detected [Fig. 3(B)]. When co-transfected with pcDNA3-



**Fig. 3** Eukaryotic expression of a novel short form of interleukin-18 (*IL-18s*)


(A) Confocal microscopy analysis of the cellular distribution of IL-18 and IL-18s in COS-7 cells. COS-7 cells were transiently transfected with myc-tagged IL-18 and IL-18s expression plasmids, and stained with mouse anti-myc monoclonal antibody and fluorescein-isothiocyanate (FITC)-labeled secondary antibody 24 h later. In these merged images, IL-18 (a) or IL-18s (b) is shown as green and nuclei as blue. The non-transfected cell as negative control is indicated by an arrow. (B) The expression of IL-18s in COS-7 cells compared with IL-18. From lane 1 to lane 5, the cells were directly lysed with loading buffer. Lane 1 contained lysates from mock transfected cells and COS-7 cells were transfected with an expression plasmid for IL-18 (lane 2), IL-18s (lane 3) alone, or in combination with the expression plasmid encoding caspase-1, respectively (lanes 4 and 5). In lanes 6 and 7, cells were transfected with IL-18s and IL-18s+caspase-1, respectively, and gently lysed followed by immunoprecipitation. Molecular mass markers are shown on the left. (C) The expression of IL-18s in media compared with IL-18. COS-7 cells were transfected with expression plasmids. Media were directly concentrated 200-fold by ultrafiltration (lanes 1–3) or concentrated 10-fold followed by immunoprecipitation (lanes 4 and 5) and analyzed by immunoblotting.

caspase-1, the mature IL-18 (approximately 18 kDa) but not the mature IL-18s, could be detected in the cell lysates [Fig. 3(B)]. When the cells were lysed in immunoprecipitation buffer and immunoprecipitated with anti-IL-18 antibody, both the pro-IL-18s and mature IL-18s could be detected [Fig. 3(B)].

For the secretion of IL-18s, the culture was analyzed by immunoblotting 72 h post-transfection. A band (approximately 16 kDa) for mature IL-18s could be detected either in concentrated preparations or in the immunoprecipitated protein preparations [Fig. 3(C)] when pcDNA3-IL-18s was co-transfected with pcDNA3-caspase-1. The expression pattern with a strong band of mature IL-18s and a faint band of proIL-18s is similar to that of IL-18 [Fig. 3(C)]. All these data suggest that caspase-1 can cleave proIL-18s and facilitate the secretion of mature IL-18s.

#### Expression, purification of GST-proIL-18s and production of mouse IL-18s

We constructed proIL-18 and proIL-18s expression vectors in which the caspase-1 cleavage site was replaced with the factor Xa site [29]. Expression of the GST-proIL-18s and GST-proIL-18 was induced by IPTG in *E. coli*. After induction, a band of approximately 43 kDa corresponding to the fusion protein was specifically bound to glutathione-Sepharose beads (Fig. 4). After extensive washing, the beads were digested with factor Xa or




**Fig. 4** Prokaryotic expression and purification of a novel short form of interleukin-18 (IL-18s) and IL-18 in mouse

*Escherichia coli* containing the expression plasmids for IL-18 and IL-18s were cultured and induced as described in "Materials and Methods". The supernatants of the bacterial sonicates were affinity-purified on glutathione agarose and samples subjected to sodium dodecyl sulphate-polyacrylamide gel electrophoresis (12%–15%). 1, molecular mass marker; 2, affinity-purified GST-proIL-18s fusion protein; 3–5, purified mouse IL-18s (16 kDa), proIL-18s (22 kDa) and IL-18 (18 kDa), respectively, after the cleavage of fusion proteins. The gel was stained with Coomassie blue.

thrombin protease. All the purified proteins were analysed by SDS-PAGE. Fig. 4 shows the purified proteins of mature IL-18s, proIL-18s and mature IL-18.

#### IL-18s lacks IL-18-like IFN- $\gamma$ -inducing activity

We evaluated whether IL-18s, like IL-18, could stimulate IFN- $\gamma$  production in mouse splenocytes. As expected, pep-IL-18 markedly stimulated IFN- $\gamma$  production in a dose-dependent manner [Fig. 5(B)], whereas the mature IL-18s did not stimulate IFN- $\gamma$  production at the concen-



**Fig. 5** Novel short form of interleukin-18 (IL-18s) enhances IL-18-induced interferon (IFN)- $\gamma$  production

(A) Mature IL-18s at 5x series dilution (0.8–500 ng/ml) was added to the spleen cells in 24-well plates alone or combined with IL-18 (16 ng/ml) and stimulated with lipopolysaccharide (LPS). After 24 h, the supernatants were collected and detected with enzyme-linked immunosorbent assay (ELISA). The control was only stimulated with LPS. The mean of three separate experiments is shown. \* $P$  < 0.05 vs. control group. (B) IL-18 at 2x series dilution (8–64 ng/ml) was added to the spleen cells in 24-well plates alone or combined with IL-18s (20 ng/ml). After 24 h, the supernatants were collected and detected with ELISA. The mean of three separate experiments is shown.

tration of 0.8–500 ng/ml (5×series dilution) [Fig. 5(A)]; similar results were obtained for proIL-18s (data not shown).

### Ability of IL-18 to induce IFN- $\gamma$ production enhanced by IL-18s

In order to determine whether IL-18s affects the activity of IL-18, the mouse splenocytes were stimulated with 2×series diluted pep-IL-18 (8–64 ng/ml) alone or combination with IL-18s (20 ng/ml). As shown in Fig. 5 (B), at low concentrations of IL-18, IL-18s increased the production of IL-18-induced IFN- $\gamma$ . At 16 ng/ml of IL-18, the production of IFN- $\gamma$  was increased from 14.9 ng/ml to 21.1 ng/ml (an approximately 40% increase) ( $P < 0.01$ ). When the concentration of IL-18 increased to 64 ng/ml, this effect was not clearly observed ( $P > 0.05$ ). Furthermore, the proIL-18s did not affect the activity of IL-18 (data not shown). These results suggest that IL-18s might enhance the activity of IL-18 when the concentration of IL-18 is comparatively low.

## Discussion

Here we reported that a short form of IL-18, termed IL-18s, 57 bp shorter than IL-18, was constitutively expressed in a wide range of mouse tissues including spleen, liver, lung, heart, brain, kidney, pancreas and gut. The expression of IL-18s was not markedly affected by LPS in splenocytes or peritoneal macrophage (data not shown).

The deleted 57 bp fragment, corresponding to nucleotide 355–411 of the full-length *IL-18* cDNA, is located at the 5' end of exon 7 of the *IL-18* gene [Fig. 1(B)]. The presence of the dinucleotide A<sup>410</sup>G<sup>411</sup> at the 3' end of the 57 bp deletion suggests that *IL-18* and *IL-18s* appear to be from the same pro-mRNA and IL-18s is an alternative splicing variant, which is in good accordance with the eukaryotic splice donor-acceptor site sequence AG/GU [23]. This dinucleotide AG is conserved not only in mouse, but also in rat and human, which might suggest that *IL-18s* exists in different species. Our results provided new evidence indicating the wide existence of *IL-18s* mRNA. We also found that, as well as a 24 kDa band corresponding to proIL-18, an approximately 21–22 kDa band could be detected in the fresh isolated splenocytes, possibly corresponding to proIL-18s. Furthermore, Pirhonen *et al.* revealed a protein (approximately 22 kDa) existed in human macrophage [33]. All of these evidences suggest that IL-18s might widely exist, and is constitutively

expressed at the levels of both mRNA and protein like IL-18.

The proIL-18 requires caspase-1 cleavage to become a biologically active molecule. Caspase-1 can enhance the maturation of proIL-18 and facilitate the export of the mature IL-18 from cells. But the caspase-1-dependent mechanism for the maturation and secretion of IL-18 has not yet been clarified, although a small amount of proIL-18 can be secreted from cells independent of caspase-1 [18–22]. In the present study, when IL-18s was transiently expressed in COS-7 cells combined with caspase-1, the mature IL-18s protein (approximately 16 kDa) could not be detected in the cell lysates [Fig. 3(B)], but could be detected in the cell lysates by immunoprecipitation and in the cell culture supernatants. We suggest that the possible reasons for IL-18s secretion might be: first, the process of caspase-1-dependent maturation and caspase-1-enhanced secretion may be closely-linked; second, the secreted mature IL-18 could bind to the surface of COS-7 cells, but the mature IL-18s could bind very weakly (data not shown), so there must be some secreted mature IL-18 counted as an intracellular part and identified in the cell lysates; third, the transient expression level of IL-18s was relatively low compared with that of IL-18; fourth, it might be a problem of recognition limitation by the anti-IL-18 antibody; and finally, the mature IL-18s detected in cell lysates with the method of immunoprecipitation might be partly due to the further digestion of caspase-1 during the process of cell lysis and immunoprecipitation.

IL-18 has a 12- $\beta$ -pleated sheet structure and an IL-1 signature-like sequence (136–160 amino acids), which is a characteristic of the IL-1 family [2,14]. For the deletion of 19 amino acids (119–137 aa), the IL-1 signature-like sequence in IL-18s is not integrated: the first conserved amino acid F136 is deleted and the eighth  $\beta$ -strand also partly deleted, which may influence the configuration of IL-18s. Furthermore, it was reported that the residues K79, E84 and D98 of mouse IL-18 are important for binding IL-18R $\beta$  to the binary complex and for biological activity [34]. In IL-18s, the residue D98 is deleted and the other two residues are very close to the 19 deleted amino acids, which suggests that the deletion in IL-18s may greatly affect the interaction with IL-18R. Consistent with this, neither proIL-18s nor mature IL-18s induced any IFN- $\gamma$  production. Although IL-18s could weakly bind to the IL-18R extracellular domain (data not shown), it might fail to recruit the IL-18R $\beta$  chain to form a functionally active ternary complex with the IL-18R $\alpha$  chain similar to that of IL-1F7b [35].

Until now, the function of IL-18s has not been clearly known. IL-18s slightly enhances the ability of IL-18 to

induce IFN- $\gamma$ , primarily at relatively low concentrations of IL-18, which might be part of its function. We presume that IL-18s might act as a naturally expressed modulator of IL-18 activity *in vivo*. Under appropriate stimuli, the concentrations of IL-18 and IL-18s increase in local microenvironments and more IFN- $\gamma$  is induced. When IL-18s reaches its working concentration, the concentration of IL-18 may be too high for IL-18s to further enhance the activities of IL-18. Increasing levels of IFN- $\gamma$  induce further expression of IL-18BPs, which slows down the increasing activity of IL-18 and gradually decreases the concentration of active IL-18. When IL-18 descends to a relatively low concentration, IL-18s might exert its function and enhance the production of IFN- $\gamma$ . To further understand the function of this novel IL-18 isoform, more researches based on these primary results are needed.

## References

- Okamura H, Tsutsui H, Komatsu T, Yutsudo M, Hakura A, Tanimoto T, Torigoe K *et al.* Cloning of a new cytokine that induces IFN- $\gamma$  production by T cells. *Nature* 1995, 378: 88–91
- Ushio S, Namba M, Okura T, Hattori K, Nukada Y, Akita K, Tanabe F *et al.* Cloning of the cDNA for human IFN-gamma-inducing factor, expression in *Escherichia coli*, and studies on the biologic activities of the protein. *J Immunol* 1996, 156: 4274–4279
- Dao T, Ohashi K, Kayano T, Kurimoto M, Okamura H. Interferon- $\gamma$ -inducing factor, a novel cytokine, enhances Fas ligand-mediated cytotoxicity of murine T helper 1 cells. *Cell Immunol* 1996, 173: 230–235
- Tsutsui H, Matsui K, Kawada N, Hyodo Y, Hayashi N, Okamura H, Higashino K *et al.* IL-18 accounts for both TNF-alpha- and Fas ligand-mediated hepatotoxic pathways in endotoxin-induced liver injury in mice. *J Immunol* 1997, 159: 3961–3967
- Dao T, Mehal WZ, Crispe IN. IL-18 augments perforin-dependent cytotoxicity of liver NK-T cells. *J Immunol* 1998, 161: 2217–2222
- Hashimoto W, Osaki T, Okamura H, Robbins PD, Kurimoto M, Nagata S, Lotze MT *et al.* Differential antitumor effects of administration of recombinant IL-18 or recombinant IL-12 are mediated primarily by Fas-Fas ligand- and perforin-induced tumor apoptosis, respectively. *J Immunol* 1999, 163: 583–589
- Hyodo Y, Matsui K, Hayashi N, Tsutsui H, Kashiwamura S, Yamauchi H, Hiroishi K *et al.* IL-18 up-regulates perforin-mediated NK activity without increasing perforin messenger RNA expression by binding to constitutively expressed IL-18 receptor. *J Immunol* 1999, 162: 1662–1668
- Takeda K, Tsutsui H, Yoshimoto T, Adachi O, Yoshida N, Kishimoto T, Okamura H *et al.* Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity* 1998, 8: 383–390
- Oshikawa K, Shi F, Rakhmievich AL, Sondel PM, Mahvi DM, Yang NS. Synergistic inhibition of tumor growth in a murine mammary adenocarcinoma model by combinational gene therapy using IL-12, pro-IL-18, and IL-1 $\beta$  converting enzyme cDNA. *Proc Natl Acad Sci USA* 1999, 96: 13351–13356
- Netea MG, Kullberg BJ, Verschuieren I, van der Meer JW. Interleukin-18 induces production of proinflammatory cytokines in mice: No intermediate role for the cytokines of the tumor necrosis factor family and interleukin-1 $\beta$ . *Eur J Immunol* 2000, 30: 3057–3060
- Hoshino T, Wiltout RH, Young HA. IL-18 is a potent coinducer of IL-13 in NK and T cells: A new potential role for IL-18 in modulating the immune response. *J Immunol* 1999, 162: 5070–5077
- Fehniger TA, Shah MH, Turner MJ, VanDeusen JB, Whitman SP, Cooper MA, Suzuki K *et al.* Differential cytokine and chemokine gene expression by human NK cells following activation with IL-18 or IL-15 in combination with IL-12: Implications for the innate immune response. *J Immunol* 1999, 162: 4511–4520
- Dinarello CA. IL-18: A TH1-inducing, proinflammatory cytokine and new member of the IL-1 family. *J Allergy Clin Immunol* 1999, 103: 11–24
- Bazan JF, Timans JC, Kastelein RA. A newly defined interleukin-1. *Nature* 1996, 379: 591
- Torigoe K, Ushio S, Okura T, Kobayashi S, Taniai M, Kunikata T, Murakami T *et al.* Purification and characterization of the human interleukin-18 receptor. *J Biol Chem* 1997, 272: 25737–25742
- Born TL, Thomassen E, Bird TA, Sims JE. Cloning of a novel receptor subunit, AcPL, required for interleukin-18 signaling. *J Biol Chem* 1998, 273: 29445–29450
- Thomassen E, Bird TA, Renshaw BR, Kennedy MK, Sims JE. Binding of interleukin-18 to the interleukin-1 receptor homologous receptor IL-1Rrp1 leads to activation of signaling pathways similar to those used by interleukin-1. *Interferon Cytokine Res* 1998, 18: 1077–1088
- Pirhonen J. Regulation of IL-18 expression in virus infection. *Scand J Immunol* 2001, 53: 533–539
- Gu Y, Kuida K, Tsutsui H, Ku G, Hsiao K, Fleming MA, Hayashi N *et al.* Activation of interferon- $\gamma$  inducing factor mediated by interleukin-1 $\beta$  converting enzyme. *Science* 1997, 275: 206–209
- Ghayur T, Banerjee S, Hugunin M, Butler D, Herzog L, Carter A, Quintal L *et al.* Caspase-1 processes IFN- $\gamma$ -inducing factor and regulates LPS-induced IFN- $\gamma$  production. *Nature* 1997, 386: 619–623
- Nakanishi K, Yoshimoto T, Tsutsui H, Okamura H. Interleukin-18 regulates both Th1 and Th2 responses. *Annu Rev Immunol* 2001, 19: 423–474
- Puren AJ, Fantuzzi G, Dinarello CA. Gene expression, synthesis, and secretion of interleukin 18 and interleukin 1 $\beta$  are differentially regulated in human blood mononuclear cells and mouse spleen cells. *Proc Natl Acad Sci USA* 1999, 96: 2256–2261
- Tone M, Thompson SA, Tone Y, Fairchild PJ, Waldmann H. Regulation of IL-18 (IFN-gamma-inducing factor) gene expression. *J Immunol* 1997, 159: 6156–6163
- Kim YM, Kang HS, Paik SG, Pyun KH, Anderson KL, Torbett BE, Choi I. Roles of IFN consensus sequence binding protein and PU.1 in regulating IL-18 gene expression. *J Immunol* 1999, 163: 2000–2007
- Gardella S, Andrei C, Poggi A, Zocchi MR, Rubartelli A. Control of interleukin-18 secretion by dendritic cells: Role of calcium influxes. *FEBS Lett* 2000, 481: 245–248
- Shida K, Shiratori I, Matsumoto M, Fukumori Y, Matsuhisa A, Kikkawa S, Tsuji S *et al.* An alternative form of IL-18 in human blood plasma: Complex formation with IgM defined by monoclonal antibodies. *J Immunol* 2001, 166: 6671–6679
- Fantuzzi G, Puren AJ, Harding MW, Livingston DJ, Dinarello CA. Interleukin-18 regulation of interferon  $\gamma$  production and cell proliferation as shown in interleukin-1 $\beta$ -converting enzyme (caspase-1)-deficient mice. *Blood* 1998, 91: 2118–2125
- Kim SH, Azam T, Yoon DY, Reznikov LL, Novick D, Rubinstein M, Dinarello CA. Site-specific mutations in the mature form of human IL-18 with enhanced biological activity and decreased neutralization by IL-18 binding protein. *Proc Natl Acad Sci USA* 2001, 98: 3304–3309
- Yang YJ, Chen SH, Ge XR. The high efficiency of expression and purification of authentic mIL-18 with high biological activity. *Curr Immunol* (forthcoming)
- Liu B, Novick D, Kim SH, Rubinstein M. Production of a biologically

- active human interleukin 18 requires its prior synthesis as PRO-IL-18. *Cytokine* 2000, 12: 1519–1525
- 31 Zou J, Bird S, Truckle J, Bols N, Horne M, Secombes C. Identification and expression analysis of an IL-18 homologue and its alternatively spliced form in rainbow trout (*Oncorhynchus mykiss*). *Eur J Biochem* 2004, 271: 1913–1923
- 32 Conti B, Jahng JW, Tinti C, Son JH, Joh TH. Induction of interferon- $\gamma$  inducing factor in the adrenal cortex. *J Biol Chem* 1997, 272: 2035–2037
- 33 Pirhonen J, Sareneva T, Julkunen I, Matikainen S. Virus infection induces proteolytic processing of IL-18 in human macrophages via caspase-1 and caspase-3 activation. *Eur J Immunol* 2001, 31: 726–733
- 34 Kato Z, Jee J, Shikano H, Mishima M, Ohki I, Ohnishi H, Li A *et al.* The structure and binding mode of interleukin-18. *Nat Struct Biol* 2003, 10: 966–971
- 35 Bufler P, Azam T, Gamboni-Robertson F, Reznikov LL, Kumar S, Dinarello CA, Kim SH. A complex of the IL-1 homologue IL-1F7b and IL-18-binding protein reduces IL-18 activity. *Proc Natl Acad Sci USA* 2002, 99: 13723–13728

Edited by  
**Lian YU**